



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

XXIV. *An Observation of an Eclipse of the Sun at the Island of New-found-land, August 5, 1766, by Mr. James Cook, with the Longitude of the Place of Observation deduced from it: Communicated by J. Bevis, M. D. F. R. S.*

Read April 30,
1767.

MR. Cook, a good mathematician, and very expert in his business, having been appointed by the Lords Commissioners of the Admiralty, to survey the sea coasts of New-found-land, Labradore, &c. took with him a very good apparatus of instruments, and among them a brass telescopic quadrant made by Mr. John Bird.

Being, August 5, 1766, at one of the Burgeo Islands near Cape Ray, latitude $47^{\circ} 36' 19''$, the south-west extremity of New-found-land, and having carefully rectified his quadrant, he waited for the eclipse of the sun; just a minute after the beginning of which, he observed the zenith distance of the sun's upper limb $31^{\circ} 57' 00''$; and, allowing for refraction and his semidiameter, the true zenith distance of the sun's centre $32^{\circ} 13' 30''$, from whence he concluded the eclipse to have begun at $0^h 4' 48''$ apparent time, and by a like process to have ended at $3^h 45' 26''$ apparent time.

N. B. There

N.B. There were three several observers, with good telescopes, who all agreed as to the moments of beginning and ending.

Mr. Cook having communicated his observation to me, I shewed it to Mr. George Witchell, who told me he had a very exact observation of the same eclipse, taken at Oxford by the Rev. Mr. Hornsby ; and he would compute, from the comparifon, the difference of longitude of the places of observation, making due allowance for the effect of parallax, and the earth's prolate spheroidal figure ; and he has since given me the following result:

5 ^h 23' 59"	beginn. at Oxford.	7 ^h 7' 5"	end at Oxford.
0 46 48	beginn. at Borgeo Ifles.	3 39 14	end at Borgeo Ifles.
<hr/>			
4 37 11		3 27 51	
— 51 59	effect of parallax, &c.	+ 17 35	effect of parallax, &c.
<hr/>			
3 45 22	diff. of meridians.	3 45 26	diff. of meridians.

J. Bevis.